

Black-Market Value of Antipsychotics, Antidepressants, and Hypnotics in Las Vegas, Nevada

TO THE EDITOR: To our knowledge, this letter represents the first effort to document the monetary value of several antipsychotic and antidepressant medications outside of a custody setting on the black-market in a major U.S. city. The monetary value of benzodiazepines and narcotics in Canada has been previously published (1), and we include a few of these medications in this letter as well. Stimulants are not included in this letter because of insufficient reports of street prices in our cohort.

Several reports have documented the potential misuse of medications, not traditionally considered to be addictive, for recreational purposes, primarily in custody settings. These include quetiapine (2, 3), anticholinergics (4), and tricyclics (5).

When publicly funded outpatient clinic staff members were asked if they had any suspicions about misuse of these medications, many said that they had heard of patients selling their medications on the street, and several suspected that some patients feigned psychosis to obtain medications to sell.

Between Oct. and Dec. of 2005, a variety of health professionals were asked if they had heard the street prices of medications. The results from sixty-one buyer and seller reports are summarized in Table 1.

TABLE 1. Results From Sixty-One Buyer and Seller Reports

Medication	Strength	Price Per Bottle (25–30 Doses)	Single Dose Price
Antipsychotics			
Olanzapine	10 mg	\$90–\$150	\$5–\$12
Quetiapine	25 mg	\$40–\$50	\$3–\$8
Antidepressants			
Mirtazapine	15 mg	\$30–\$43	\$3–\$5
Citalopram	10 mg	\$20–\$30	
Fluoxetine	20 mg	\$22–\$29	
Hypnotics			
Clonazepam	1 mg	\$50–\$80	\$5–\$11
Diazepam	5 mg	\$35–\$60	\$5–\$7
Zolpidem	5 mg	\$20–\$28	\$3–\$6
Alprazolam	0.5 mg	\$12–\$22	

Sellers reported that it was relatively easy to sell the medications they had received free of charge from the clinics and convert them into money for rent, utilities, food, illicit drugs, or alcohol. Buyers reported that they use these medications for their sedative effect as a sleep aid, to “zone-out” or to “take the edge off.” Buyers included individuals attempting to self-medicate, not having a third-party method of payment.

More detailed studies should be performed to elucidate the abuse potential of these and other medications as well as the epidemiology of their misuse. The health effects of black-market use are as yet unstudied, and physicians should be aware that patients might be covertly taking psychotropic medicines. Diversion of these medications may represent a significant expense for public and private agencies, and suspected malingering or other requests for early refills should alert agencies to the possibility of black-market activity.

References

1. Sajan A, Corneil T, Grzybowski S: The street value of prescription drugs. *CMAJ* 1998; 159:139–142
2. Pierre JM, Shnyder I, Wirshing DA, Wirshing WC: Intranasal quetiapine abuse (letter). *Am J Psychiatry* 2004; 161:1718
3. Hussain MZ, Waheed W, Hussain S: Intravenous quetiapine abuse (letter). *Am J Psychiatry* 2005; 162:1755–1756
4. Buhrich N, Weller A, Kevans P: Misuse of anticholinergic drugs by people with serious mental illness. *Psychiatr Serv* 2000; 51: 928–929
5. Hepburn S, Harden J, Grieve JHK, Hiscox J: Deliberate misuse of tricyclic antidepressants by intravenous drug users: case studies and report. *Scott Med J* 2005; 50:131–133

GREG TARASOFF, M.D.
KATHRYN OSTI
Las Vegas, Nev.

Physical Exercise as a Treatment for Non-Suicidal Self-Injury: Evidence From a Single-Case Study

TO THE EDITOR: Nonsuicidal self-injury refers to deliberate damage to one's own bodily tissue without suicidal intent. Nonsuicidal self-injury is a pervasive behavior problem, yet effective treatments are lacking. Research has found that people engage in nonsuicidal self-injury for several different reasons, including tension relief and the induction of pleasurable affective states (1). Preliminary evidence suggests that the release of endogenous opioids may be central to this process of emotional regulation (2). We hypothesized that aerobic physical exercise, which has been shown to regulate mood (3) and stimulate the release of beta-endorphin (4), would decrease the frequency of nonsuicidal self-injury. We report on the results of a single-case study demonstrating the effectiveness of physical exercise as a treatment for nonsuicidal self-injury.

“Ms. A” was an overweight 26-year-old woman with a 13-year history of ongoing psychological and pharmacological treatment for persistent nonsuicidal self-injury, including one inpatient hospitalization for nonsuicidal self-injury within the past year. Ms. A was receiving twice-weekly outpatient psychotherapy for the duration of this study. In an initial baseline assessment, she reported 2.25 episodes of nonsuicidal self-injury per week over the previous month, including self-hitting and head-banging.

We provided Ms. A with a 60-minute workout video, instructing her to exercise three times per week and to exercise in response to nonsuicidal self-injury urges at any time. We also provided a daily assessment form in which she recorded mood and self-injurious urges (both on 0–9 scales) and behaviors.

The frequency of Ms. A's nonsuicidal self-injury decreased immediately after the introduction of exercise to 0.37 times per week during a five-week experimental phase. She then independently discontinued exercise. During this quasi-experimental return to baseline, nonsuicidal self-injury increased to 2.33 times per week. When exercise was re-introduced, nonsuicidal self-injury decreased to 0.00 times per week for the remainder of the study.

Overall, nonsuicidal self-injury frequency was significantly lower during exercise phases ($M=0.29$, $SD=0.49$) relative to nonexercise phases ($[M=2.20$, $SD=0.45]$ $t=6.93$, $df=10$, $p<0.001$). Moreover, analysis of mood ratings showed an increase from before exercise ($M=2.23$, $SD=$